

## E.1 DISSOCIATION CONSTANTS FOR ACIDS AT 25 °C

Name	Formula	$K_{a1}$	$K_{a2}$	$K_{a3}$
Acetic	$\text{HC}_2\text{H}_3\text{O}_2$	$1.8 \times 10^{-5}$		
Ascorbic	$\text{HC}_6\text{H}_7\text{O}_6$	$8.0 \times 10^{-3}$		
Arsenic	$\text{H}_3\text{AsO}_4$	$5.6 \times 10^{-3}$	$1.0 \times 10^{-7}$	$3.0 \times 10^{-12}$
Arsenous	$\text{H}_3\text{AsO}_3$	$6.0 \times 10^{-10}$		
Benzoic	$\text{HC}_7\text{H}_5\text{O}_2$	$6.5 \times 10^{-5}$		
Boric	$\text{H}_3\text{BO}_3$	$5.8 \times 10^{-10}$		
Butyric acid	$\text{HC}_4\text{H}_7\text{O}_2$	$1.5 \times 10^{-5}$		
Carbonic	$\text{H}_2\text{CO}_3$	$4.3 \times 10^{-7}$	$5.6 \times 10^{-11}$	
Chloroacetic	$\text{HC}_2\text{H}_2\text{O}_2\text{Cl}$	$1.4 \times 10^{-3}$		
Cyanic	$\text{HCNO}$	$3.5 \times 10^{-4}$		
Citric	$\text{H}_3\text{C}_6\text{H}_5\text{O}_7$	$7.4 \times 10^{-4}$	$1.7 \times 10^{-5}$	$4.0 \times 10^{-7}$
Hydroazotic	$\text{HN}_3$	$1.9 \times 10^{-5}$		
Hydrocyanic	$\text{HCN}$	$4.9 \times 10^{-10}$		
Hydrogen chromate ion	$\text{HCrO}_4^-$	$3.0 \times 10^{-7}$		
Hydrogen peroxide	$\text{H}_2\text{O}_2$	$2.4 \times 10^{-12}$		
Hydrogen selenate ion	$\text{HSeO}_4^-$	$2.2 \times 10^{-2}$		
Hydrogen sulfate ion	$\text{HSO}_4^-$	$1.2 \times 10^{-2}$		
Hydrogen sulfide	$\text{H}_2\text{S}$	$5.7 \times 10^{-8}$	$1.3 \times 10^{-13}$	
Hypobromous	$\text{HBrO}$	$2.0 \times 10^{-9}$		
Hypochlorous	$\text{HClO}$	$3.0 \times 10^{-8}$		
Hypoiodous	$\text{HIO}$	$2.0 \times 10^{-11}$		
Iodic	$\text{HIO}_3$	$1.7 \times 10^{-1}$		
Lactic	$\text{HC}_3\text{H}_5\text{O}_3$	$1.4 \times 10^{-4}$		
Malonic	$\text{H}_2\text{C}_3\text{H}_2\text{O}_4$	$1.5 \times 10^{-3}$	$2.0 \times 10^{-6}$	
Oxalic	$\text{H}_2\text{C}_2\text{O}_4$	$5.9 \times 10^{-2}$	$6.4 \times 10^{-5}$	
Phenol	$\text{HC}_6\text{H}_5\text{O}$	$1.3 \times 10^{-10}$		
Phosphoric	$\text{H}_3\text{PO}_4$	$7.5 \times 10^{-3}$	$6.2 \times 10^{-8}$	$4.2 \times 10^{-13}$
Paraperiodic	$\text{H}_5\text{IO}_6$	$2.8 \times 10^{-2}$	$5.3 \times 10^{-9}$	
Propionic	$\text{HC}_3\text{H}_5\text{O}_2$	$1.3 \times 10^{-5}$		
Pyrophosphoric	$\text{H}_4\text{P}_2\text{O}_7$	$3.0 \times 10^{-2}$	$4.4 \times 10^{-3}$	
Selenous	$\text{H}_2\text{SeO}_3$	$2.3 \times 10^{-3}$	$5.3 \times 10^{-9}$	
Sulfuric	$\text{H}_2\text{SO}_4$	strong acid	$1.2 \times 10^{-2}$	
Sulfurous	$\text{H}_2\text{SO}_3$	$1.7 \times 10^{-2}$	$6.4 \times 10^{-8}$	
Tartaric	$\text{H}_2\text{C}_4\text{H}_4\text{O}_6$	$1.0 \times 10^{-3}$	$4.6 \times 10^{-5}$	

## E.2 DISSOCIATION CONSTANTS FOR BASES AT 25 °C

Name	Formula	$K_b$	Name	Formula	$K_b$
Ammonia	$\text{NH}_3$	$1.8 \times 10^{-5}$	Hydroxylamine	$\text{HONH}_2$	$1.1 \times 10^{-8}$
Aniline	$\text{C}_6\text{H}_5\text{NH}_2$	$4.3 \times 10^{-10}$	Methylamine	$\text{CH}_3\text{NH}_2$	$4.4 \times 10^{-4}$
Dimethylamine	$(\text{CH}_3)_2\text{NH}$	$5.4 \times 10^{-4}$	Pyridine	$\text{C}_5\text{H}_5\text{N}$	$1.7 \times 10^{-9}$
Ethylamine	$\text{C}_2\text{H}_5\text{NH}_2$	$6.4 \times 10^{-4}$	Trimethylamine	$(\text{CH}_3)_3\text{N}$	$6.4 \times 10^{-5}$
Hydrazine	$\text{H}_2\text{NNH}_2$	$1.3 \times 10^{-6}$			